

In the Claims

The claims are as follows:

1. (Previously Presented) An isolated synthetic peptide or polypeptide comprising a domain which specifically binds a nucleic acid sequence linked to a domain which specifically binds a metal which is hydrolytically or redox active linked to a further domain which specifically binds a nucleic acid sequence, wherein the domain which binds a metal is capable of binding a lanthanide, calcium or a metal in the same group as calcium.
2. (Previously Presented) The peptide or polypeptide of claim 1 which comprises the amino acid sequence TERRRQQLDKDGDGTIDEREIKIHFQNKRAKIK (SEQ ID NO:2), or a portion thereof which binds the nucleic acid sequence and the metal.
3. (Withdrawn) The peptide or polypeptide of claim 1 which comprises the amino acid sequence TERRRFDKDGNGYISAAELRHVKIWFQNKRAKIK (SEQ ID NO:3), or a catalytically active portion thereof.
4. (Original) The peptide or polypeptide of claim 1 which comprises at least 20 residues.
5. (Original) The peptide or polypeptide of claim 1 which comprises a consensus EF-Hand sequence.
6. (Previously Presented) The peptide or polypeptide of claim 1 wherein one domain which specifically binds a nucleic acid sequence is a domain from a transcription factor.
7. (Previously Presented) The peptide or polypeptide of claim 1 wherein one domain which specifically binds a nucleic acid sequence comprises a helix or a strand from a helix-turn-helix motif, a relaxed helix-turn-helix motif, a winged helix-turn-helix motif, a helix-loop-strand motif, or a hormone receptor motif.

8. (Original) The peptide or polypeptide of claim 1 which binds dsDNA, dsRNA, ssDNA, ssRNA, A-DNA, B-DNA, or Z-DNA.
9. (Original) The peptide or polypeptide of claim 6 wherein the transcription factor is engrailed.
10. (Previously Presented) The peptide or polypeptide of claim 1 wherein the domains which specifically bind nucleic acid are from a homeodomain.
11. (Original) The peptide or polypeptide of claim 6 wherein the transcription factor comprises a helix-turn-helix domain.
12. (Original) The peptide or polypeptide of claim 7 which comprises alpha-helices 2 and 3 of a helix-turn-helix motif.
- 13-15. (Canceled)
16. (Original) The peptide or polypeptide of claim 1 which binds Ca(II) or Mg(II).
17. (Original) The peptide or polypeptide of claim 1 which binds Eu(III).
18. (Original) The peptide or polypeptide of claim 1 further comprising a protein transport domain.
19. (Withdrawn) An isolated nucleic acid molecule comprising a nucleic acid segment which encodes the peptide or polypeptide of claim 1, or a sequence complementary thereto.
20. (Withdrawn) An expression cassette comprising the nucleic acid molecule of claim 19 which is operably linked to a promoter functional in a host cell.

-
21. (Withdrawn) A host cell comprising the expression cassette of claim 20.
 22. (Withdrawn) The host cell of claim 21 wherein the host cell is a prokaryotic cell.
 23. (Withdrawn) The host cell of claim 21 wherein the host cell is a eukaryotic cell.
 24. (Withdrawn) The host cell of claim 21 wherein the host cell is a plant cell.
 25. (Withdrawn) A vector comprising the expression of cassette of claim 20.
 26. (Withdrawn) A method to identify a synthetic endonuclease comprising:
 - a) identifying a secondary structure in a first amino acid sequence that binds to a specific nucleic acid sequence which has geometrical similarity to the secondary structure of a second amino acid sequence that binds a metal; and
 - b) identifying a third amino acid sequence that comprises at least a portion of the first amino acid sequence and at least a portion of the second amino acid sequence to yield a synthetic endonuclease which specifically binds the specific nucleic acid sequence and specifically binds the metal.
 27. (Original) A method of using a synthetic endonuclease specific for a particular nucleic acid sequence comprising: contacting a sample comprising isolated nucleic acid with an amount of the peptide or polypeptide of claim 1 effective to cleave at least one nucleic acid sequence in the sample.
 28. (Original) A method to detect the presence of a nucleic acid sequence in a sample comprising:
 - a) contacting a sample comprising nucleic acid suspected of containing a nucleic acid sequence recognized by the peptide or polypeptide of claim 1, with the peptide or polypeptide of claim 1; and

- b) determining or detecting whether the peptide or polypeptide cleaves the nucleic acid sequence.
29. (Original) The method of claim 28 wherein the detection or determination is by gel electrophoresis.
30. (Withdrawn) The peptide or polypeptide of claim 1 which comprises TERRRFRVFDKDGNGYISAAEKIWFQNKRAKIK (SEQ ID NO:4), or a catalytically active portion thereof.
31. (Withdrawn) The peptide or polypeptide of claim 1 which comprises the amino acid sequence TRRRRFLSFDKDGDTITTKEEVWFQNRMRKWK (SEQ ID NO:5), or a catalytically active portion thereof.
32. (Withdrawn) The peptide or polypeptide of claim 1 which comprises the amino acid sequence DEKRPRTAFSGEQLARLKREFNENRYLTERRRLRVFDKDGNGF ISAAEKI WFQNK RAKIKKST (SEQ ID NO:6), or a catalytically active portion thereof.
33. (Withdrawn) The peptide or polypeptide of claim 1 which comprises the amino acid sequence TERRRQQLDKDGDGTIDEREIKIWFQNKRAKIK (SEQ ID NO:7), or a catalytically active portion thereof.
34. (Previously Presented) The peptide or polypeptide of claim 1 which binds a lanthanide (III) or (IV).
35. (Previously Presented) The peptide or polypeptide of claim 1 which comprises the amino acid sequence FRVFDKDGNGYISAAEKIWF (SEQ ID NO:8), or a catalytically active portion thereof.

SUPPLEMENTAL AMENDMENT AND EXAMINER INTERVIEW SUMMARY

Serial Number: 09/785,546

Filing Date: February 16, 2001

Title: ARTIFICIAL ENDONUCLEASE

Page 6

Dkt: 875.037US1

36. (Previously Presented) The peptide or polypeptide of claim 1 wherein the lanthanide is cerium, praseodymium, neodymium, promethium, samarium, europium, gadolinium, terbium, dysprosium, holmium, erbium, thulium, ytterbium, or lutetium.